



0.33 fl oz (10mL) bottle NDC 35781-0210-1  
1.01 fl oz (30mL) bottle NDC 35781-0210-2

## READ THIS INFORMATION BEFORE PRESCRIBING THIS PRODUCT

### DESCRIPTION OF SpeedGel Rx®

SpeedGel Rx® is a homeopathic topical analgesic gel that contains the ingredients listed below. Homeopathic ingredients have been used since the inception of this science and remain as an effective method of treating select conditions. It is an amber colored odorless gel for use externally to control inflammation and reduce pain.

#### It contains the following **active ingredients**:

Aconitum Napellus 3X HPUS, Arnica Montana 1X HPUS, Belladonna 3X HPUS, Bellis Perennis 1X HPUS, Calendula Officinalis 1X HPUS, Chamomilla 1X HPUS, Colchicum 3X HPUS, Echinacea Angustifolia 1X HPUS, Echinacea Purpurea 1X HPUS, Hamamelis Virginiana 1X HPUS, Hypericum Perforatum 1X HPUS, Millefolium 1X HPUS, Symphytum Officinale 3X HPUS, Zingiber Officinale 1X HPUS.

#### It also contains the following **inactive ingredients**:

Docusate sodium, Isopropyl myristate, Lecithin, Purified water, Sodium hydroxide, Urea.

### CLINICAL PHARMACOLOGY

The exact pharmacology by which SpeedGel Rx® works to control aches and pains associated with arthritis or trauma (such as sprains, strains, dislocations, repetitive/overuse injuries, traumatic edema, post surgical edema, hematoma, general swelling of joints and soft tissues) to such areas as hands, wrist, elbow, shoulder, neck, back, knees, ankles, feet and toes is unknown. It is theorized that, according to traditional homeopathy, the following ingredients contribute in the following ways:

Ingredients	Common Name	Indications
Aconitum Napellus	Monk's-hood	Neuralgia, rheumatism, hemostasis, analgesia
Arnica Montana	Mountain arnica	Stimulates healing of injured tissues, wounds, contusions, hematomas, neuralgia, myalgia, analgesia
Belladonna	Deadly nightshade	Inflammation locally
Bellis Perennis	Daisy	Dislocations, bruising, reducing edema
Calendula Officinalis	Calendula	Analgesia, inflammation
Colchicum	Colchicine	Inflammation and gout
Chamomilla	Chamomile	Inflammation, promotes healing of tissues
Echinacea Angustifolia	Narrow leaf cone flower	Inflammation and inhibits hyaluronidase
Echinacea Purpurea	Purple cone flower	Stimulates fibroblasts, inflammation
Hamamelis Virginiana	Witch-hazel	Astringent, analgesic, hematomas
Hypericum Perforatum	St. John's wort	Neuropathic pains
Millefolium	Yarrow	Hematomas, wound healing
Symphytum Officinale	Comfrey	Neuropathy, causalgia, contusions, periostitis
Zingiber Officinale	Ginger	Inflammation

### INDICATIONS AND USAGE

SpeedGel Rx® is indicated for the relief of pain and inflammation from: arthritis or injuries such as sprains, strains, dislocations, repetitive/overuse injuries, traumatic edema, post surgical edema, general swelling of joints and soft tissues of the hand, wrist, elbow, shoulder, neck, back, knee, ankle, foot and toe and general aches and pains.

### CONTRAINDICATIONS

Known sensitivity to Arnica montana.

### WARNINGS

For external use only. Direct patient not to ingest SpeedGel Rx® and to avoid contact with the eyes and mucous membranes, wounds, and damaged skin. If condition worsens, or if symptoms persist for more than seven days or clear up and occur again within a few days, patient should consult a doctor. If a rash develops, patient should discontinue use until rash clears. After the disappearance of rash, patient can try SpeedGel Rx® again on a test area and monitor the site for additional results. If no rash or redness results, then patient can resume use. However, if the rash persists or redevelops, use should be discontinued.

Direct patient to keep this product out of reach of children and seek medical help or contact a Poison Control Center immediately if swallowed.

### PRECAUTIONS

Use in Pregnancy: No human or animal studies on the effect of SpeedGel Rx® in pregnancy have been conducted. Nursing Mothers: It is not known whether SpeedGel Rx® is excreted in breast milk.

### ADVERSE REACTIONS

Rarely, allergic skin reactions may occur. These effects are transient and will clear after a few days.

### DOSAGE AND ADMINISTRATION

Apply a thin layer (1 drop covers an area of skin 2 inches by 2 inches) to the affected area 3-4 times over a 24-hour period, or as directed by a physician, and rub in gently. Applications of less than 3-4 times a day will not produce optimum results. Excess drops may be wiped from the area. Safe to use on children over age of 2 years.

### HOW SUPPLIED

Package Size: 1.01 fl oz (30mL) bottle

Manufactured for:  
Gensco Laboratories, LLC  
Miami, FL 33122

866 608 6284  
855 7GENSCO  
855 743 6726

[www.speedgelrx.com](http://www.speedgelrx.com)